
SADLIER

ProgressMathematics

Standards-Based Instruction & Practice



Aligned to the

Common Core State Standards for Mathematics

Grade 7

Contents

Ratios & Proportional Relationships	4
The Number System	1
Expressions & Equations	4
Geometry	
Statistics & Probability	6



Ratios & Proportional Relationships

7.RP

GRADE 7 MATHEMATICS STANDARDS		SADLIER PROGRESS MATHEMATICS, GRADE 7
Analyze proportional relationships and use them to solve real-world and mathematical problems.		
CCSS.MATH.CONTENT.7.RP.A.1 Compute unit rates associated with ratios of fractions, including ratios of lengths, areas and other quantities measured in like or different units. For example, if a person walks 1/2 mile in each 1/4 hour, compute the unit rate as the complex fraction 1/2/1/4 miles per hour, equivalently 2 miles per hour.	Lesson 1	Compute Unit Rates—pp. 10-17
CCSS.MATH.CONTENT.7.RP.A.2 Recognize and represent proportional relationships between quantities.		
CCSS.MATH.CONTENT.7.RP.A.2.A Decide whether two quantities are in a proportional relationship, e.g., by testing for equivalent ratios in a table or graphing on a coordinate plane and observing whether the graph is a straight line through the origin.	Lesson 2	Identify Proportional Relationships—pp. 18– 25
CCSS.MATH.CONTENT.7.RP.A.2.B Identify the constant of proportionality (unit rate) in tables, graphs, equations, diagrams, and verbal descriptions of proportional relationships.	Lesson 3	Identify the Constant of Proportionality —pp. 26–33
CCSS.MATH.CONTENT.7.RP.A.2.C Represent proportional relationships by equations. For example, if total cost t is proportional to the number n of items purchased at a constant price p , the relationship between the total cost and the number of items can be expressed as $t = pn$.	Lesson 4	Represent Proportional Relationships with Equations —pp. 34–41
CCSS.MATH.CONTENT.7.RP.A.2.D Explain what a point (x, y) on the graph of a proportional relationship means in terms of the situation, with special attention to the points (0, 0) and (1, r) where r is the unit rate.	Lesson 5	Interpret Graphs of Proportional Relationships—pp. 42–49
CCSS.MATH.CONTENT.7.RP.A.3 Use proportional relationships to solve multistep ratio and percent problems. Examples: simple interest, tax, markups and markdowns, gratuities and commissions, fees, percent increase and decrease, percent error.	Lesson 6	Problem Solving: Multi-step Ratio Problems— pp. 50–57
	Lesson 7	Problem Solving: Multi-step Percent Problems—pp. 58–65

The Number System

7.NS

GRADE 7 MATHEMATICS STANDARDS		SADLIER PROGRESS MATHEMATICS, GRADE 7
Apply and extend previous understandings of operations with fractions to add, subtract, multiply, and divide rational numbers.		
CCSS.MATH.CONTENT.7.NS.A.1 Apply and extend previous understandings of addition and subtraction to add and subtract rational numbers; represent addition and subtraction on a horizontal or vertical number line diagram.		
CCSS.MATH.CONTENT.7.NS.A.1.A Describe situations in which opposite quantities combine to make 0. For example, a hydrogen atom has 0 charge because its two constituents are oppositely charged.	Lesson 8	Understand Addition of Integers—pp. 72-79
CCSS.MATH.CONTENT.7.NS.A.1.B Understand $p + q$ as the number located a distance $ q $ from p , in the positive or negative direction depending on whether q is positive or negative. Show that a number and its opposite have a sum of 0 (are additive inverses). Interpret sums of rational numbers by describing real-world contexts.	Lesson 8	Understand Addition of Integers—pp. 72-79
CCSS.MATH.CONTENT.7.NS.A.1.C Understand subtraction of rational numbers as adding the additive inverse, $p - q = p + (-q)$. Show that the distance between two rational numbers on the number line is the absolute value of their difference, and apply this principle in real-world contexts.	Lesson 9	Understand Subtraction of Integers—pp. 80–87
CCSS.MATH.CONTENT.7.NS.A.1.D Apply properties of operations as strategies to add and subtract rational numbers.	Lesson 10	Add and Subtract Rational Numbers —pp. 88–95
CCSS.MATH.CONTENT.7.NS.A.2 Apply and extend previous understandings of multiplication and division and of fractions to multiply and divide rational numbers.		
CCSS.MATH.CONTENT.7.NS.A.2.A Understand that multiplication is extended from fractions to rational numbers by requiring that operations continue to satisfy the properties of operations, particularly the distributive property, leading to products such as (-1)(-1) = 1 and the rules for multiplying signed numbers. Interpret products of rational numbers by describing real-world contexts.	Lesson 8	Understand Addition of Integers—pp. 72-79
CCSS.MATH.CONTENT.7.NS.A.2.B Understand that integers can be divided, provided that the divisor is not zero, and every quotient of integers (with non-zero divisor) is a rational number. If p and q are integers, then $-(p/q) = (-p)/q = p/(-q)$. Interpret quotients of rational numbers by describing real-world contexts.	Lesson 12	Understand Division of Integers—pp. 104–111

The Number System

7.NS

GRADE 7 MATHEMATICS STANDARDS		SADLIER PROGRESS MATHEMATICS, GRADE 7
CCSS.MATH.CONTENT.7.NS.A.2.C Apply properties of operations as strategies to multiply and divide rational numbers.	Lesson 13	Multiply and Divide Rational Numbers—pp. 112–119
CCSS.MATH.CONTENT.7.NS.A.2.D Convert a rational number to a decimal using long division; know that the decimal form of a rational number terminates in 0s or eventually repeats.	Lesson 14	Convert Rational Numbers to Decimal Form— pp. 120–127
CCSS.MATH.CONTENT.7.RP.A.1 Solve real-world and mathematical problems involving the four operations with rational numbers. 1.0. **The content of the conte	Lesson 15	Apply Rational-Number Operations —pp. 128- 135
¹ Computations with rational numbers extend the rules for manipulating fractions to complex fractions.		

Expressions & Equations		7.EE
GRADE 7 MATHEMATICS STANDARDS		SADLIER PROGRESS MATHEMATICS, GRADE 7
Use properties of operations to generate equivalent expressions.		
CCSS.MATH.CONTENT.7.EE.A.1 Apply properties of operations as strategies to add, subtract,	Lesson 16	Combine Like Terms to Simplify Linear Expressions —pp. 142–149
factor, and expand linear expressions with rational coefficients.	Lesson 17	Expand and Factor Linear Expressions—pp. 150–157
CCSS.MATH.CONTENT.7.EE.A.2 Understand that rewriting an expression in different forms in a	Lesson 16	Combine Like Terms to Simplify Linear Expressions — pp. 142–149
problem context can shed light on the problem and how the quantities in it are related. For example, $a + 0.05a = 1.05a$ means that "increase by 5%" is the same as "multiply by 1.05."	Lesson 17	Expand and Factor Linear Expressions—pp. 150–157
Solve real-life and mathematical problems using numerical and algebraic expressions and equations.		
CCSS.MATH.CONTENT.7.EE.B.3 Solve multi-step real-life and mathematical problems posed with positive and negative rational numbers in any form (whole numbers, fractions, and decimals), using tools strategically. Apply properties of operations to calculate with numbers in any form; convert between forms as appropriate; and assess the reasonableness of answers using mental computation and estimation strategies. For example: If a woman making \$25 an hour gets a 10% raise, she will make an additional 1/10 of her salary an hour, or \$2.50, for a new salary of \$27.50. If you want to place a towel bar 9 3/4 inches long in the center of a door that is 27 1/2 inches wide, you will need to place the bar about 9 inches from each edge; this estimate can be used as a check on the exact computation.	Lesson 18	Problem Solving: Multi-step Problems with Rational Numbers —pp. 158–165

Expressions & Equations

7.EE

Grade 7 Mathematics Standards		SADLIER PROGRESS MATHEMATICS, GRADE 7
CCSS.MATH.CONTENT.7.EE.B.4 Use variables to represent quantities in a real-world or mathematical problem, and construct simple equations and inequalities to solve problems by reasoning about the quantities.		
CCSS.MATH.CONTENT.7.EE.B.4.A	Lesson 19	Solve Linear Equations—pp. 166–173
Solve word problems leading to equations of the form $px + q = r$ and $p(x + q) = r$, where p , q , and r are specific rational numbers. Solve equations of these forms fluently. Compare an algebraic solution to an arithmetic solution, identifying the sequence of the operations used in each approach. For example, the perimeter of a rectangle is 54 cm. Its length is 6 cm. What is its width?	Lesson 20	Problem Solving: Linear Equations—pp. 174– 181
CCSS.MATH.CONTENT.7.EE.B.4.B	Lesson 21	Solve Linear Inequalities—pp. 182-189
Solve word problems leading to inequalities of the form $px + q > r$ or $px + q < r$, where p , q , and r are specific rational numbers. Graph the solution set of the inequality and interpret it in the context of the problem. For example: As a salesperson, you are paid \$50 per week plus \$3 per sale. This week you want your pay to be at least \$100. Write an inequality for the number of sales you need to make, and describe the solutions.	Lesson 22	Problem Solving: Linear Inequalities—pp. 190–197

Geometry 7.G

Grade 7 Mathematics Standards		SADLIER PROGRESS MATHEMATICS, GRADE 7
Draw, construct, and describe geometrical figures and describe the relationships between them.		
CCSS.MATH.CONTENT.7.G.A.1 Solve problems involving scale drawings of geometric figures, including computing actual lengths and areas from a scale drawing and reproducing a scale drawing at a different scale.	Lesson 23	Use Scale Drawings to Solve Problems —p 204–211
CCSS.MATH.CONTENT.7.G.A.2 Draw (freehand, with ruler and protractor, and with	Lesson 24	Draw Shapes that Meet Given Conditions -212-219
technology) geometric shapes with given conditions. Focus on constructing triangles from three measures of angles or sides, noticing when the conditions determine a unique triangle, more than one triangle, or no triangle.	Lesson 25	Construct Triangles Using Both Side Lengt and Angle Measures—pp. 220–227
CCSS.MATH.CONTENT.7.G.A.3 Describe the two-dimensional figures that result from slicing three-dimensional figures, as in plane sections of right rectangular prisms and right rectangular pyramids.	Lesson 26	Slice Three-Dimensional Figures—pp. 228-

Geometry	7.G
----------	-----

GRADE 7 MATHEMATICS STANDARDS		SADLIER PROGRESS MATHEMATICS, GRADE 7
Solve real-life and mathematical problems involving angle measure, area, surface area, and volume.		
CCSS.MATH.CONTENT.7.G.B.4 Know the formulas for the area and circumference of a circle and use them to solve problems; give an informal derivation of the relationship between the circumference and area of a circle.	Lesson 27	Use Formulas for Area and Circumference of Circles—pp. 236–243
CCSS.MATH.CONTENT.7.G.B.5 Use facts about supplementary, complementary, vertical, and adjacent angles in a multi-step problem to write and solve simple equations for an unknown angle in a figure.	Lesson 28	Use Equations to Find Unknown Angle Measures—pp. 244–251
CCSS.MATH.CONTENT.7.G.B.6 Solve real-world and mathematical problems involving area, volume and surface area of two- and three-dimensional objects composed of triangles, quadrilaterals, polygons, cubes, and right prisms.	Lesson 29	Problem Solving: Area, Volume, and Surface Area—pp. 252–259
Statistics & Probability		7.SP
GRADE 7 MATHEMATICS STANDARDS		SADLIER PROGRESS MATHEMATICS, GRADE 7
Use random sampling to draw inferences about a population.		
CCSS.MATH.CONTENT.7.SP.A.1 Understand that statistics can be used to gain information about a population by examining a sample of the population; generalizations about a population from a sample are valid only if the sample is representative of that population. Understand that random sampling tends to produce representative samples and support valid inferences.	Lesson 30	Understand Sampling—pp. 266–273
Use data from a random sample to draw inferences about a population with an unknown characteristic of interest. Generate multiple samples (or simulated samples) of the same size to gauge the variation in estimates or predictions. For example, estimate the mean word length in a book by randomly sampling words from the book; predict the winner of a school election based on randomly sampled survey data. Gauge how far off the estimate or prediction might be.	Lesson 31	Use Sampling to Draw Inferences—pp. 274–281
Draw informal comparative inferences about two populations.		
CCSS.MATH.CONTENT.7.SP.B.3 Informally assess the degree of visual overlap of two numerical data distributions with similar variabilities, — continued —	Lesson 32	Use Visual Overlap to Compare Distributions—pp. 282–289

Statistics & Probability

7.SP

GRADE 7 MATHEMATICS STANDARDS		SADLIER PROGRESS MATHEMATICS, GRADE 7
as a multiple of a measure of variability. For example, the mean height of players on the basketball team is 10 cm greater than the mean height of players on the soccer team, about twice the variability (mean absolute deviation) on either team; on a dot plot, the separation between the two distributions of heights is noticeable.		
CCSS.MATH.CONTENT.7.SP.B.4 Use measures of center and measures of variability for numerical data from random samples to draw informal comparative inferences about two populations. For example, decide whether the words in a chapter of a seventh-grade science book are generally longer than the words in a chapter of a fourth-grade science book.	Lesson 33	Use Sample Statistics to Compare Populations—pp. 290–297
Investigate chance processes and develop, use, and evaluate probability models.		
CCSS.MATH.CONTENT.7.SP.C.5 Understand that the probability of a chance event is a number between 0 and 1 that expresses the likelihood of the event occurring. Larger numbers indicate greater likelihood. A probability near 0 indicates an unlikely event, a probability around 1/2 indicates an event that is neither unlikely nor likely, and a probability near 1 indicates a likely event.	Lesson 34	Understand Probability of a Chance Event—pp. 298–305
CCSS.MATH.CONTENT.7.SP.C.6 Approximate the probability of a chance event by collecting data on the chance process that produces it and observing its long-run relative frequency, and predict the approximate relative frequency given the probability. For example, when rolling a number cube 600 times, predict that a 3 or 6 would be rolled roughly 200 times, but probably not exactly 200 times.	Lesson 35	Relate Relative Frequency and Probability—pp. 306–313
CCSS.MATH.CONTENT.7.SP.C.7 Develop a probability model and use it to find probabilities of events. Compare probabilities from a model to observed frequencies; if the agreement is not good, explain possible sources of the discrepancy.		
CCSS.MATH.CONTENT.7.SP.C.7.A Develop a uniform probability model by assigning equal probability to all outcomes, and use the model to determine probabilities of events. For example, if a student is selected at random from a class, find the probability that Jane will be selected and the probability that a girl will be selected.	Lesson 36	Develop a Uniform Probability Model —pp. 314–321
CCSS.MATH.CONTENT.7.SP.C.7.B Develop a probability model (which may not be uniform) by observing frequencies in data generated from a chance process. For example, find the approximate probability that a spinning penny will land heads up or that a tossed paper cup will land open-end down. Do the outcomes for the spinning penny appear to be equally likely based on the observed frequencies?	Lesson 40	Summarize Numerical Data—pp. 346–353

Statistics & Probability

7.SP

Grade 7 Mathematics Standards		SADLIER PROGRESS MATHEMATICS, GRADE 7
CCSS.MATH.CONTENT.7.SP.C.8 Find probabilities of compound events using organized lists, tables, tree diagrams, and simulation.		
CCSS.MATH.CONTENT.7.SP.C.8.A Understand that, just as with simple events, the probability of a compound event is the fraction of outcomes in the sample space for which the compound event occurs.	Lesson 38	Find Probabilities of Compound Events—pp. 330–337
CCSS.MATH.CONTENT.7.SP.C.8.B Represent sample spaces for compound events using methods such as organized lists, tables and tree diagrams. For an event described in everyday language (e.g., "rolling double sixes"), identify the outcomes in the sample space which compose the event.	Lesson 39	Represent Sample Spaces for Compound Events—pp. 338–345
CCSS.MATH.CONTENT.7.SP.C.8.C Design and use a simulation to generate frequencies for compound events. For example, use random digits as a simulation tool to approximate the answer to the question: If 40% of donors have type A blood, what is the probability that it will take at least 4 donors to find one with type A blood?	Lesson 40	Simulate Compound Events—pp. 346–353